

Dimerization of atomic chains

Andriy Zhugayevych (zhugayevych@iop.kiev.ua)

November 20, 2011

1	Introduction	2
2	Electronic states of dimerized chain	3
2.1	Solving the Schroedinger equation	3
2.2	Perfectly dimerized chain	3
2.3	Band energy and density matrix	5
3	Electronic properties of defects	6
3.1	Calculating localized states	6
3.2	Calculating transmission amplitude	7
3.3	Analytic properties of the transmission amplitude	9
3.4	Topological defects in hopping rate dimerization	10
3.5	Topological defects in on-site energy dimerization	11
3.6	Pair of t-kinks	11
3.7	Pair of e- and t-kinks	13
4	Electron-electron interaction	13
4.1	Hartree-Fock method in ZDO approximation	14
4.2	Three-diagonal approximation	14
4.3	Coulomb potential in three-diagonal approximation	15
4.4	Spin contamination: SUHF	16
4.5	Perfectly dimerized chain: renormalization equations	17
4.6	RHF solutions in absence of “bare” dimerization	19
4.7	UHF solutions in absence of “bare” dimerization	21
4.8	“Bare” t-dimerization	21
5	Electron-phonon interaction	22
5.1	Electron-phonon interaction in adiabatic approximation	22
5.2	Effects of dielectric environment	23
5.3	Hartree–Fock description	25
5.4	Extended SSH model: infinite chain	26
5.5	Extended SSH model: infinite chain (old version)	27
5.6	Hubbard–Peierls model	28
6	Phonons	30
7	Continuous model of dimerized chain	30
7.1	Basic equations	30
7.2	Topological defects	32
7.3	Kinks	33
7.4	Kink-antikink pair	34
7.5	Continuous model	34
7.6	Nonlinear excitations	35
7.7	Nonlinear excitations: effects of electron-electron interaction	37
7.8	Jackiw–Rebbi model	38
8	Conjugated polymers	39
8.1	Model parameters for polyacetylene	39
8.2	Can we think about single and double bonds in polyacetylene?	41
	Appendix	41
A	Evaluation of transmission amplitude	41
B	RHF for perfectly dimerized chain in the most general setup	41
C	Derivation of the electronic Hamiltonian	42
D	Formulas for the gradient and Hessian	43
E	Participation ratio	44
	References	44